

## HEDIS® Provider Guide:

### Measure Description

Children 3 months to 18 years of age diagnosed with Upper Respiratory Infection (URI) **should not** be dispensed an antibiotic within 3 days of the diagnosis.

Note: Claims/encounters with more than one diagnosis (e.g., competing diagnoses) are excluded from the measure.

### Using Correct Billing Codes

#### Codes to Identify URI

Description	ICD-9 Code	ICD-10 Code*
Acute Nasopharyngitis (common cold)	460	J00
Acute Laryngopharyngitis	465.0	J06.0
Acute URI	465.8, 465.9	J06.0

\* ICD-10 -CM codes are to be used on or after 10/1/2015.

#### Codes to Identify Competing Diagnoses

Description	ICD-9 Code	ICD-10 Code*
Otitis Media	382	H66, H67
Acute Sinusitis	461	J01.80, J01.90
Chronic Sinusitis	473	J32
Pharyngitis	462	J02.0, J02.8, J02.9

\* ICD-10 -CM codes are to be used on or after 10/1/2015.

Description	ICD-9 Code	ICD-10 Code*
Streptococcal Tonsillitis	034.0	J03.00, J03.01, J03.80
Acute Tonsillitis	463	J03.81, J03.90, J03.91
Pneumonia	418-486	J13-J20

## How to Improve HEDIS® Scores

- **Do not** prescribe an antibiotic for a URI diagnosis only.
- Submit any co-morbid/competing diagnosis codes that apply (examples listed above in the 'Codes to Identify Common Competing Diagnoses').
- Code and bill for all diagnoses based on patient assessment.
- Educate patient on comfort measures (e.g., acetaminophen for fever, rest, extra fluids) and advise patient to call back if symptoms worsen (antibiotic can be prescribed if necessary after 3 days of initial diagnosis).
- You are encouraged to resubmit an encounter if you missed a second diagnosis code and you see a member on the non-compliant list in Cozeva®.
- Patient education materials on antibiotic resistance and common infections can be found here: <http://www.cdc.gov/getsmart/index.html>