

HEDIS® Provider Guide:

Measure Description

Women 16-24 years of age who were identified as sexually active and who had at least one Chlamydia test during the measurement year.

Members identified as sexually active:

- Pregnancy
- STI screening
- Contraceptive prescribed
- Pregnancy test completed

Exclusions:

- Pregnancy test and a prescription for isotretinoin within 6 days
- Pregnancy test and an x-ray within 6 days

Using Correct Billing Codes

Description	CPT Code
Chlamydia Screening	87110, 87270, 87320, 87490-87492, 87810

Contraceptives

<ul style="list-style-type: none"> • Desogestrel-ethinyl estradiol • Dienogest-estradiol multiphasic • Drospirenone-ethinyl estradiol • Drospirenone-ethinyl estradiol-levomefolate biphasic • Ethinyl estradiol-ethynodiol • Ethinyl estradiol-etonogestrel • Ethinyl estradiol-folic acid-levonorgestrel • Ethinyl estradiol-levonorgestrel 	<ul style="list-style-type: none"> • Ethinyl estradiol-norelgestromin • Ethinyl estradiol-norethindrone • Ethinyl estradiol-norgestimate • Ethinyl estradiol-norgestrel • Etonogestrel • Levonorgestrel • Medroxyprogesterone • Mestranol-norethindrone • Norethindrone
---	--

How to Improve HEDIS® Scores

- Perform Chlamydia screening every year on every 16-24-year old female identified as sexually active (use any visit opportunity).
- Add Chlamydia screening as a standard lab for women 16-24 years old. Use adolescent well-care exams and well woman exams for this purpose.
- Ensure that you have an opportunity to speak with your adolescent female patient without her parents present.
- Remember that Chlamydia screening can be performed through a urine test. Offer this option for your patients.